



Reumatismo - The Italian Journal of Rheumatology

<https://www.reumatismo.org/reuma>

eISSN 2240-2683

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The Early Access service lets users access peer-reviewed articles well before print/regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

Reumatismo is, therefore, E-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination, and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

The E-publishing of this PDF file has been approved by the authors.

Please cite this article as:

Reis do Espírito Santos R, Muniz Caldas CA, Freire de Carvalho J. **Peripheral neuropathy in antiphospholipid syndrome: a systematic review.** *Reumatismo* doi: 10.4081/reumatismo.2025.1735

Submitted: 05-04-2024

Accepted: 26-09-2024

 © the Author(s), 2025
Licensee PAGEPress, Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Peripheral neuropathy in antiphospholipid syndrome: a systematic review

Rafael Reis do Espírito Santos,¹ Cezar Augusto Muniz Caldas,¹ Jozélio Freire de Carvalho²

¹Faculty of Medicine, Federal University of Pará, Belém-Pará; ²Institute for Health Sciences, Federal University of Bahia, Salvador, Brazil

Correspondence: Jozélio Freire de Carvalho, Institute for Health Sciences, Federal University of Bahia, Salvador, Brazil

E-mail: jotafc@gmail.com

Key words: peripheral neuropathy, antiphospholipid syndrome, thrombosis, thrombophilia, mononeuritis multiplex

Contributions: CAC, writing, data analysis, review; JFC, design, data collection, writing, data analysis, statistical analysis, submission; RBES, data analysis, writing, revision.

Conflict of interest: the authors declare no potential conflict of interest.

Ethics approval and consent to participate: the authors declare that Helsinki's World Medical Association Declaration was followed.

Informed consent: not applicable.

Availability of data and materials: all data are available upon request.

Funding: no funding or sponsorship was received for this study or publication of this article.

Summary

Objective. Antiphospholipid syndrome (APS) is a disease characterized by recurrent thrombosis in the presence of antiphospholipid antibodies. The most uncommon events described in the literature have been peripheral neurological disorders. This paper aims to systematically review the cases of peripheral neuropathy (PN) in APS patients.

Methods. We systematically searched articles on PN and APS with English abstracts in PubMed from 1966 to August 2022.

Results. We found 10 articles on PN and APS with 100 patients. Age varied from 25 to 78 years; 86-100% of patients in these studies were female. Most patients had primary APS (n=9); one article considered secondary APS associated with other autoimmune diseases. Disease duration varied from 0 to 8.6 years, but three articles did not provide this information. Most studies showed positivity for anticardiolipin antibodies (n=5), followed by lupus anticoagulant (n=2).

Regarding clinical NP features, mononeuritis multiplex (n=3) and autonomic neuropathy (n=3) were more common than peripheral polyneuropathy (n=2). Nerve biopsy was performed in 7 articles and found positivity in all cases. Concerning treatment, most articles used anticoagulation (n=4), followed by glucocorticoids (n=3), intravenous immunoglobulin, and immunosuppressive drugs (n=1). Most cases improved after treatment (n=7).

Conclusions. This study demonstrates that PN is a rare complication in APS and occurs more frequently in females, associated with antiphospholipid antibody positivity. Most cases were confirmed by electroneurography or nerve biopsy and had a good outcome.

Introduction

Antiphospholipid syndrome (APS) is an acquired thrombophilia characterized by thrombosis and/or obstetric events in the presence of persistent antiphospholipid antibodies (1). It is pathogenically mediated by antibodies targeting phospholipid-binding proteins, mainly β -2-glycoprotein I (2).

The most common manifestations are thrombotic events such as deep venous thrombosis and stroke. Additional disease presentations include limb ischemia, myocardial infarction, and pulmonary thromboembolism. The clinical spectrum of APS can also include nonvascular manifestations such as thrombocytopenia, livedo reticularis, migraine, and seizures, to name some. Microvascular thrombosis is also part of the APS and includes obstetrical events, kidney lesions, and catastrophic APS. The neurological system, mainly the central nervous system, is involved in 1 to 20% of APS cases (3). Also some peripheral neurological disorders have been described, although uncommonly (4).

Recently, a new classification for APS was developed and published by the American College of Rheumatology and the European Alliance of Rheumatologic Associations (5). The main novelty introduced by this set of criteria consists in the identification of six clinical domains (macrovascular venous thromboembolism, macrovascular arterial thrombosis, microvascular, obstetric, cardiac valve, and hematologic) and two laboratory domains (lupus anticoagulant functional coagulation assays, and solid-phase enzyme-linked immunosorbent assays for IgG/IgM anticardiolipin and/or IgG/IgM anti- β 2-glycoprotein I antibodies) (5).

This article aims to perform a systematic review of peripheral neuropathy (PN) in APS patients.

Methods

Literature review

We have systematically studied articles published in PubMed/MEDLINE, Web of Sciences, LILACS, and SciELO from 1966 to August 2022 using the following MeSH entry terms: “peripheral neuropathy” and “antiphospholipid syndrome”, without language restriction. The reference lists of the selected articles were analyzed to identify other publications. Initially, three independent investigators (JFC, CAMC, and RRES) performed the literature search and independently selected the study abstracts. In the second stage, the same reviewers independently read the full-text articles selected by abstracts. The same third reviewer also resolved disagreements arising during a consensus meeting. The authors followed the PRISMA guidelines (6). We designed a standardized form to extract the following information from relevant articles regarding the authors and year of publication, the number of patients studied, demographic data, disease duration, nerve biopsy, electroneuromyography features, treatments, and outcomes.

Results

We found ten articles on APS and PN totaling 100 patients. Table 1 summarizes these studies (4, 7-15). Most of them were case reports (n=5), with also retrospective (n=3), case-control (n=1) and cross-sectional (n=1) studies. Patients were mainly female (86-100% of the study cohort across different studies), and their ages varied from 25 to 78 years. Nine studies included patients with primary APS and one study recruited patients with secondary APS. Disease duration ranged from 0 to 8.6 years, although most papers did not provide this detail. References number 7 and 10 from the same author reported 15 patients in 2014 and 22 in 2017: it is not clear if there is an overlap between these patients.

Most studies showed positivity for anticardiolipin antibodies IgG (25/51, 50%) and IgM (35/51, 67%), followed by lupus anticoagulant (10/51, 20%), whereas three studies (n=49 patients) did not fully characterize the aPL profile.

Regarding clinical PN features, autonomic neuropathy (30/43, 69%), mononeuritis multiplex (3/43, 7%), simplex mononeuropathy (1/43, 2%), and peripheral polyneuropathy (2/43, n=2, 5%).

Nerve biopsy was performed in seven articles and found histologic abnormalities suggestive of PN in all cases.

Only five articles described the treatment for the PN, and all of them being single-case reports: most patients used anticoagulation (n=3), glucocorticoids (n=3), intravenous immunoglobulins, and immunosuppressive drugs (n=1). All the patients improved after treatment, but the outcome was not described in three articles.

Discussion

This study is the first systematic review of PN in APS.

Although the pathogenesis of PN is poorly understood, small fiber dysfunction can result from micro thrombosis, direct antibody binding to neuronal epitopes that lead to nerve dysfunction, or both mechanisms. The latter hypothesis is supported by reported symptom improvement with immune system modulatory therapy (16, 17).

The involvement of the central nervous system in APS is well known. However, the peripheral neurological manifestations are less described in the literature, with mononeuritis multiplex being the most common manifestation in 60% of our studies. The main nerves involved were sural, fibular, and median, with reports of associated carpal tunnel syndrome (7-15).

In 90% of the reported studies, the clinical diagnosis was complemented by electroneuromyography or skin biopsy. About 30% (n=3) confirmed the presence of PN using the combined method of nerve biopsy and electroneuromyography. In comparison, 60% (n=6) underwent one and the other investigation, while the electroneuromyographic study, despite being widespread, fails to indicate the cause.

Skin biopsy, on the other hand, is easy to perform and is emerging as the gold standard for diagnosing sensory and autonomic neuropathies of small fibers with an estimated sensitivity and specificity of around 90% (10), while the electroneuromyographic study despite being widespread, it fails to perform histological analysis or indicate the cause.

In the studies evaluated, 40% (n=4) used anticoagulants to treat thrombotic symptoms, thus improving the symptoms of PN (7-15). One of the studies showed improved symptoms when using warfarin orally during the 2-year follow-up, without recurrence of the disease (15). In contrast, 20% (n=2) needed methylprednisolone, followed by 20% (n=2) of intravenous immunoglobulin.

Finally, some limitations were observed in the published studies, including the fact that the number of participants is still low. Future studies should involve large patient samples with more long-term follow-up, enabling a better understanding of this neurological manifestation in APS. The studies must address exclusively primary APS, since associated diseases such as SLP may induce PN. It is known that 7.6% of lupus patients have this neurological abnormality (18). Studies using standard methodologies to detect PN, such as electroneurography, are also desired, as well as the use of nerve biopsy and standardized methods for evaluating antiphospholipid antibodies.

This study's strengths include considering studies with patients diagnosed according to international criteria for APS (19, 20).

Conclusions

The present study was a systematic review of all studies that evaluated PN in APS patients. Although uncommon, PN in APS should be considered, mainly in female patients with antiphospholipid antibody positivity. Most cases were confirmed by electroneurography or nerve biopsy and had a good outcome.

References

1. Sammaritano LR. Antiphospholipid syndrome. *Best Pract Res Clin Rheumatol* 2020; 34: 101463.
2. Levine SR, Salowich-Palm L, Sawaya KL, Perry M, Spencer HJ, et al. IgG anticardiolipin antibody titer > 40 GPL and the risk of subsequent thrombo-occlusive events and death: a prospective cohort study. *Stroke* 1997; 28: 1660-5.
3. Rodrigues CE, Carvalho JF, Shoenfeld Y. Neurological manifestations of antiphospholipid syndrome. *Eur J Clin Invest* 2010; 40: 350-9.
4. Santos MS, de Carvalho JF, Brotto M, Bonfa E, Rocha FA. Peripheral neuropathy in patients with primary antiphospholipid (Hughes') syndrome. *Lupus* 2010; 19: 583-90.
5. Barbhaiya M, Zuily S, Naden R, Hendry A, Manneville F, Amigo MC, et al. 2023 ACR/EULAR antiphospholipid syndrome classification criteria. *Ann Rheum Dis* 2023; 82: 1258-70.
6. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ* 2021; 372: n71.
7. Schofield JR. Autonomic neuropathy-in its many guises-as the initial manifestation of the antiphospholipid syndrome. *Immunol Res* 2017; 65: 532-42.
8. Giorli E, Franciotta D, Serventi A, Binda GA, Canepa G, Siciliano G, et al. Sacral nerve stimulation for fecal incontinence in a patient with antiphospholipid syndrome-related autonomic neuropathy. *Muscle Nerve* 2016; 53: 992-4.
9. Takahashi M, Katada F, Sato S, Shibayama H, Fukutake T, Murayama S. Mononeuritis multiplex due to thrombotic ischemia of primary antiphospholipid antibody syndrome without vasculitis: an autopsy case report. *Rinsho Shinkeigaku* 2015; 55: 897-903. [Article in Japanese].
10. Schofield JR, Blitshteyn S, Shoenfeld Y, Hughes GR. Postural tachycardia syndrome (POTS) and other autonomic disorders in antiphospholipid (Hughes) syndrome (APS). *Lupus* 2014; 23: 697-702.
11. Etemadifar M, Dehghani L, Tahani S, Toghianifar N, Rahaimi M, Eskandari N. Neurological manifestations in patients with antiphospholipid syndrome. *Iran J Neurol* 2013; 12: 172-5.
12. Bilora F, Biasiolo M, Zancan A, Zanon E, Veronese F, Manca F, Sartori MT. Autonomic dysfunction and primary antiphospholipid syndrome: a frequent and frightening correlation? *Int J Gen Med* 2012; 5: 339-43.
13. Rodrigues CE, Carvalho JF. Multiple mononeuropathy secondary to thrombosis of the vasa nervorum in primary antiphospholipid syndrome. *Isr Med Assoc J* 2010; 12: 448-9.
14. Jeruc J, Popovic M, Vizjak A, Jurčić V, Lestan B, Ferluga D. Multiple mononeuropathy due to vasculitis associated with anticardiolipin antibodies: a case report. *Folia Neuropathol* 2006; 44: 140-3.
15. Erten N, Saka B, Karan MA, Parman Y, Umman B, Tascioglu C. Catastrophic secondary antiphospholipid syndrome with peripheral nervous system involvement: a case report. *Acta Med Okayama* 2004; 58: 107-10.
16. Sanna G, D'Cruz D, Cuadrado MJ. Cerebral manifestations in the antiphospholipid (Hughes) syndrome. *Rheum Dis Clin North Am* 2006; 32: 465-90.
17. Katzav A, Chapman J, Shoenfeld Y. CNS dysfunction in the antiphospholipid syndrome. *Lupus* 2003; 12: 903-7.
18. Hanly JG, Li Q, Su L, Urowitz MB, Gordon C, Bae SC, et al. Peripheral nervous system disease in systemic lupus erythematosus: results from an international inception cohort study. *Arthritis Rheumatol* 2020; 72: 67-77.
19. Wilson WA, Gharavi AE, Koike T, Lockshin MD, Branch DW, Piette JC, et al. International consensus statement on preliminary classification criteria for definite antiphospholipid syndrome: report of an international workshop. *Arthritis Rheum* 1999; 42: 1309-11.
20. Miyakis S, Lockshin MD, Atsumi T, Branch DW, Brey RL, Cervera R, et al. International consensus statement on an update of the classification criteria for definite antiphospholipid syndrome (APS). *J Thromb Haemost* 2006; 4: 295-306.

Table 1. Summary of the studies on antiphospholipid syndrome and peripheral neuropathy.

Author, year,	Study design	N, female sex	Age, years	Primary or secondary APS	Disease duration	APS clinical manifestations	aPL antibodies	PN alterations	ENMG	Nerve biopsy	Treatment	Outcome
Schofield, 2017 [7]	Retrospective	22, 86%	26.5 (6-48)	Primary	4.4 years	59% thrombotic events	IgM aCL 95%	Autonomic neuropathy (POTS, NCS)	ND	Autonomic neuropathy by skin biopsy	ND	ND
Giorli <i>et al.</i> , 2016 [8]	Case report	1, 100%	54	Primary	ND	ND	ND	weakness in distal muscles in all limbs and reduced tendon stretch reflexes in the lower limbs without sensory deficits	Moderately severe polyneuropathy	Mild inflammatory demyelinating neuropathy	High dose GC and IVIg-no response; plasmapheresis and azathioprine; neurostimulation	Improved muscle strenght.
Takahashi <i>et al.</i> , 2015 [9]	Autopsy case report	1, male	78	Primary	ND	Livedo, purpura	ND	Mononeuritis multiplex	ND	Axonal degeneration and thrombi on arterioles and venules, without inflammation	Anticoagulation	He improved of PN. But, 2 months later died from venous embolism and infection
Schofield <i>et al.</i> , 2014 [10]	Retrospective	15, 93%	39 (15-66)	Primary and secondary (SLE 2, Sjogren 1, RA 1, Crohn 1, celiac disease 1)		Livedo, Raynaud phenomenon,	ND	Postural tachycardia syndrome (N=8), neurocardiogenic syncope (n=8), and orthostatic hypotension	ND	Positive skin biopsy	ND	Improved
Etemadifar <i>et al.</i> , 2013 [11]	Retrospective	1/103, 87%	36.3 ±11.4	Primary	ND	DVT, PTE	ND	ND	ND	ND	ND	ND
Bilora <i>et al.</i> , 2012 [12]	Case-control	31 APS, 31 controls	47.4 ±21	Primary	ND	ND	ND	ND	ND	ND	ND	ND
Santos <i>et al.</i> , 2010 [4]	Cross-sectional	26, 92%	38.6 ±8.5	Primary	8.6 ± 3	Arterial: 59%; Venous: 50%; Obstetric:41%	LA: 41%; IgG aCL: 86%; IgM aCL: 55%	Mononeuritis multiplex	41% altered	ND	ND	ND
Rodrigues <i>et al.</i> , 2010 [13]	Case report	1, 100%	25	Primary	0	Cyanosis of the lower limbs, livedo, malleolar ulcers.	54 GPL, 28 MPL, LA	Pain and partial loss of flexion and extension of the right ankle	Marked degeneration of motor and sensory nere fibers of the posterior tibial and fibular nerves, sensorial of the sural compatible wit moneurite multiplex.	Fibrin thrombus of the vasa nervosum, without inflammation	Methylprednisolone pulse therapy and anticoagulation	Improved, complete recovery in 11 months
Jeruc <i>et al.</i> , 2006 [14]	Case report	1, 100%	49	Primary	ND	Abortion, Raynaud's phenomenon	IgG aCL	Multiple mononeuropathy	ND	Active necrotizing arteritis of the epineural arteries with transmural inflammatory infiltrate and thrombosis	Methylprednisolone 48mg <i>per ors</i>	Improved
Erten <i>et al.</i> , 2004 Turkey [15]	Case report	1, 100%	34	Primary	ND	CAPS (heart failure, stroke cerebral venous thrombosis, multiple cranial and abdominal thromboses)	aCL 27GPL	Peripheral polyneuropathy	Mononeuropathy multiplex with severe axonal degeneration	Severe axonal degeneration with perivascular inflammation related to wallerian degeneration caused by ischemy.	Anticoagulation	Improved. Complete recovery.

aCL, anticardiolipin; aPL, antiphospholipid antibodies; APS, antiphospholipid syndrome; CAPS, catastrophic antiphospholipid syndrome; ENMG, electroneuromyography; GC, glucocorticoid; IVIg, intravenous immunoglobulin; LA, lupus anticoagulant; N, number; ND, not described; PN, peripheral neuropathy; POTS, postural orthostatic tachycardia syndrome; NCS, neurocardiogenic syncope.